

## **DEPARTMENT OF HEALTH & HUMAN SERVICES**

**Public Health Service** 

Centers for Disease Control and Prevention

September 21, 2017

Anjani Chandra, Ph.D.
Principal Investigator and Team Leader
National Survey of Family Growth
National Center for Health Statistics
3311 Toledo Road
Hyattsville, MD 20782

Dear Dr. Chandra,

I am writing in support of the National Survey of Family Growth (NSFG), and to emphasize its importance in monitoring HIV prevention activities. The Division of HIV/AIDS Prevention (DHAP) at the Centers for Disease Control and Prevention (CDC) is responsible for funding HIV prevention activities in the United States and evaluating those efforts. Although much of what we know about HIV infection has come from HIV/AIDS case surveillance data and from behavioral surveillance studies of groups at increased risk for acquiring HIV or people living with HIV, data from national probability samples such as the NSFG are also critical to our ability to monitor HIV in the United States.

The NSFG is a source of several important indicators we use to monitor progress toward national goals. These include indicators of sexual risk such as multiple sex partners, sex with persons who inject drugs or HIV-infected persons, and the exchange of sex for money or drugs. Each of these behaviors place individuals at increased risk for HIV infection. The NSFG also provides important information about behaviors that can reduce HIV acquisition or transmission in the U.S. population, such as condom use and HIV testing. HIV testing data provided by NSFG help evaluate the uptake of CDC's recommendations on routine HIV testing in all healthcare settings for patients between the ages of 13 through 64, as well as repeat screening every year for patients at high risk for HIV infection. Additionally, NSFG serves as the primary data source for monitoring several HIV-related objectives in Healthy People 2020, such as increasing the proportion of adolescents and adults who have ever been tested for HIV.

DHAP also uses data from NSFG to monitor indicators in populations at high risk for HIV, such as men who have sex with men (MSM), a population that accounted for 70% of all diagnoses in 2015. DHAP has frequently relied on NSFG data to analyze and

disseminate information about HIV testing and risk behaviors among MSM as well as other populations at increased risk (see references below). Furthermore, NSFG data were instrumental in enabling DHAP to produce national population size estimates for populations at risk for HIV, including MSM (Purcell 2011), persons who inject drugs (Lansky 2013), and heterosexuals at risk (Lansky 2015), which have improved our estimation of HIV rates in these populations.

The NSFG is one of only a few nationally representative surveys that collect data related to HIV risk and prevention on a regular basis. It has a history of successfully collecting sensitive information from respondents, and it continues to be an important data source for DHAP.

We look forward to continued success of the NSFG.

Sincerely yours,

Marc Pitasi, MPH

Epidemiologist

Special Studies and Diagnostics Team

Behavioral and Clinical Surveillance Branch

Division of HIV/AIDS Prevention

Centers for Disease Control and Prevention

1600 Clifton Rd. NE, MS-E46

Atlanta, GA 30333 Phone: 404 639 6361

Email: mpitasi@cdc.gov

cc: Joseph Prejean, Chief, Behavioral and Clinical Surveillance Branch

Kevin Delaney, Acting Team Lead, Special Studies and Diagnostics Team

## References

Purcell DW, Johnson CH, Lansky A, Prejean J, Stein R, et al. (2012) Estimating the population size of men who have sex with men in the United States to obtain HIV and syphilis rates. *Open AIDS J* 6: 98–107

Lansky A, Finlayson T, Johnson C, Holtzman D, Wejnert C, et al. (2014) Estimating the Number of Persons Who Inject Drugs in the United States by Meta-Analysis to Calculate National Rates of HIV and Hepatitis C Virus Infections. *PLoS ONE* 9(5): e97596

Lansky A, Johnson C, Oraka E, Sionean C, Joyce MP, et al. (2015) Estimating the Number of Heterosexual Persons in the United States to Calculate National Rates of HIV Infection. *PLoS ONE* 10(7): e0133543

Abara WE, Oraka E, Jeffries WL 4th, Chavez P, Nasrullah M, et al. (2017) Prevalence and Correlates of Condom Use Among Sexually Active Men Who Have Sex with Men in the United States: Findings from the National Survey of Family Growth, 2002, 2006–10 and 2011–13. *Sex Health* epub ahead of print. doi: 10.1071/SH16034

Aholou TM, McCree DH, Oraka E, Jeffries WL 4th, Rose CE, et al. (2017) Sexual Risk and Protective Behaviors Among Reproductive-Aged Women in the United States. *J Womens Health (Larchmt)* epub ahead of print. doi: 10.1089/jwh.2016.6224

Balaji AB, Oraka E, Fasula AM, Jayne PE, Carry MG, et al. (2017) Association Between Parent-Adolescent Communication About Sex-Related Topics and HIV Testing, United States. 2006–2013. *AIDS Care* 29(3): 344-9

Fasula AM, Oraka E, Jeffries WL 4th, Carry M, Banez Ocfemia MC, et al. (2016) Young Sexual Minority Males in the United States: Sociodemographic Characteristics and Sexual Attraction, Identity and Behavior. *Perspect Sex Reprod Health* 48(1): 3-8

Jeffries WL 4th, Johnson OD. (2015) Homonegative Attitudes and Risk Behaviors for HIV and Other Sexually Transmitted Infections Among Sexually Active Men in the United States. *Am J Public Health* 105(12): 2466-72

Nasrullah M, Oraka E, Chavez P, Valverde E, DiNenno E. (2015) Nonvolitional Sex and HIV-related Sexual Risk Behaviours Among MSM in the United States. *AIDS* 29(13): 1673-80

Nasrullah M, Oraka E, Chavez PR, Johnson CH, DiNenno E. (2017) Factors Associated with Condom Use Among Sexually Active US Adults, National Survey of Family Growth, 2006–2010 and 2011–2013. *J Sex Med* 14(4): 541-50

Van Handel M, Lyons B, Oraka E, Nasrullah M, DiNenno E, et al. (2015) Factors Associated with Time Since Last HIV Test Among Persons at High Risk for HIV Infection, National Survey of Family Growth, 2006–2010. *AIDS Patient Care STDS* 29(10): 533-40