



## Document Details

<b>Docket ID:</b>	FS-2018-0035
<b>Docket Title:</b>	Agency Information Collection Activities; Proposals, Submissions, and Approvals: Community Forest and Open Space Program
<b>Document File:</b>	
<b>Docket Phase:</b>	Notice
<b>Phase Sequence:</b>	1
<b>Original Document ID:</b>	FS-2018-0035-DRAFT-0005
<b>Current Document ID:</b>	FS-2018-0035-0006
<b>Title:</b>	Comment on FR Doc # 2018-13261
<b>Number of Attachments:</b>	0
<b>Document Type:</b>	PUBLIC SUBMISSIONS
<b>Comment on Document ID:</b>	FS-2018-0035-0001
<b>Comment on Document Title:</b>	Agency Information Collection Activities; Proposals, Submissions, and Approvals: Community Forest and Open Space Program
<b>Status:</b>	Posted
<b>Received Date:</b>	07/10/2018
<b>Date Posted:</b>	07/11/2018
<b>Posting Restriction:</b>	No restrictions
<b>Submission Type:</b>	Web
<b>Number of Submissions:</b>	1

## Document Optional Details

<b>Status Set Date:</b>	07/11/2018
<b>Current Assignee:</b>	NA
<b>Status Set By:</b>	Migliori, Michael (FS)
<b>Tracking Number:</b>	1k2-947a-bd0z
<b>Page Count:</b>	1
<b>Total Page Count Including Attachments:</b>	1

## Submitter Info

**Comment:**

Large wildfires have a considerable impact on the atmospheric concentrations of CO<sub>2</sub>, CO, O<sub>3</sub>, NO<sub>x</sub>, and CH<sub>4</sub> across North America. Carbon releases can be as high as 4 to 8 kg C-m<sup>-2</sup> per fire event. These emissions significantly affect concentrations far downwind. With funding from NASA, the Joint Fire Science Program, NSF, and the Canadian Government, US and Canadian researchers have been developing a uniform approach to estimate carbon, trace gas, and particulate emissions from wildfire. Models to estimate the consumption of the forest floor and peatlands in boreal North American are being developed to study terrestrial carbon cycling and estimate trace gas emissions. Measurements of atmospheric CO, O<sub>3</sub>, nitrogen oxides, and equivalent black carbon show boreal fires to be of great importance to levels of these gas and aerosol species on the continental to hemispheric scale. The work presented examines how fuel consumption varies in the deep surface organic layers found in many boreal forests and peatlands, and how factors controlling the variations in fuel consumption influence inter-annual to decadal variations in the North American terrestrial carbon budget and the emissions of CO, and ozone precursors. Our federal forests, managed by the Forest Service and Bureau of Land Management (BLM), urgently need active management to reduce the risk of severe wildfire. At least 58 million acres of national forest are at high or very high risk of severe wildfire, and over 4.5 million homes are at risk. Over 1.1 million acres of national forest need reforestation. But last year the U.S. Forest Service treated less than 204,000 acres, a small fraction of what's needed. A significant part of the problem is process and paperwork. It typically takes 18 months to four years for federal agencies to develop and implement forest projects. Forest Service employees typically spend 40 percent of their time doing paperwork instead of managing forests. In the West, this year's wildfire season has been one of the worst on record. Portland and Seattle have both been covered in smoke for days on end, with ash falling in the streets, schools cancelled, children huddled inside, and health-sensitive individuals suffering distress. The more than 40,000-acre Eagle Creek Fire devastated many treasured recreational sites in the Columbia River Gorge and closed a key Interstate highway for weeks. Across the country, nearly 4.5 million homes are at risk from wildfire. Catastrophic fires are the result of decades of fire suppression, coupled with unprecedented fuel buildups due to a lack of forest management activity. These catastrophic fires destroy valuable timber resources but also degrade many of the other uses of healthy forests. In one 2014 fire, nearly 20,000 acres of high-quality northern spotted owl habitat burned. In fact, over the past two decades, wildfire has become the greatest source of habitat loss for the northern spotted owl. Between 1995 and 2015, according to the Forest Service, habitat impact attributed to wildfire was ten times the impact from timber harvest. Since 2015, wildfire impacts have only worsened. One recent study showed that probability of extirpation of California spotted owls increases by a factor of seven after a severe fire. Some deny the fire science because it conflicts with their ideology. They deny that these fires are actually catastrophic, or they point to climate change to deny that fuel buildup plays any role in fire intensification. Climate change is certainly NOT a factor. Overstocked, stressed, kindling-like forests create firestorms

that outpace anything the country has seen in living memory. It is no coincidence that over 90% of the burned acres in Oregon this year were on Forest Service lands which comprise just over 50% of Oregons forestland and where active management is nearly at a standstill. The state and federal government have about equal amounts of land in Oregon, and experience equal numbers of fire starts. But burned areas are overwhelmingly concentrated on Forest Service lands. Active management will make these federal forests more resilient to these extreme events. Environmental activist must not get the way of Forest Service. Sell timber, build Sawmills, \*🌐

**First Name:** Anonymous

**Last Name:** Anonymous

**ZIP/Postal Code:**

**Email Address:**

**Government Agency Type:** 🌐

**Government Agency:** 🌐

**Category:** 🌐

**Cover Page:** 