



Department of Civil and Environmental Engineering

February 3, 2022

Docket Number: EPA-HQ-OAR-2021-0324

Comments of Drs. Fatemeh Kazemiparkouhi¹, David MacIntosh¹, Helen Suh²

¹ Environmental Health and Engineering, Inc., Newton, MA and the Harvard T.H. Chan School of Public Health, Boston, MA

² Tufts University, Medford, MA

We are writing to comment on issues raised by the proposed RFS annual rule, the Draft Regulatory Impact Analysis (December 2021; EPA-420-D-21-002), and the supporting Health Effects Docket Memo (September 21, 2021; EPA-HQ-OAR-2021-0324-0124), specifically regarding the impact of ethanol-blended fuels on air quality and public health. We provide evidence of the air quality and public health benefits provided by higher ethanol blends, as shown in our recently published study¹ by Kazemiparkouhi et al. (2021), which characterized emissions from light duty vehicles for market-based fuels. Findings from our study demonstrate ethanol-associated reductions in emissions of primary particulate matter (PM), nitrogen oxides (NOx), carbon monoxide (CO), and to a lesser extent total hydrocarbons (THC). Our results provide further evidence of the potential for ethanol-blended fuels to improve air quality and public health, particularly for environmental justice communities. Below we present RFS-pertinent findings from Kazemiparkouhi et al. (2021), followed by their implications for air quality, health, and environmental justice.

Summary of Kazemiparkouhi et al. (2021)

Our paper is the first large-scale analysis of data from light-duty vehicle emissions studies to examine real-world impacts of ethanol-blended fuels on regulated air pollutant emissions, including PM, NOx, CO, and THC. To do so, we extracted data from a comprehensive set of emissions and market fuel studies conducted in the US. Using these data, we (1) estimated composition of market fuels for different ethanol volumes and (2) developed regression models to estimate the impact of changes in ethanol volumes in market fuels on air pollutant emissions for different engine types and operating conditions. Importantly, our models estimated these changes accounting for not only ethanol volume fraction, but also aromatics volume fraction, 90% volume distillation temperature (T90) and Reid Vapor Pressure (RVP). Further, they did so under both cold start and hot stabilized running conditions and for gasoline-direct

¹ <https://doi.org/10.1016/j.scitotenv.2021.151426>

injection engines (GDI) and port-fuel injection (PFI) engine types. Key highlights from our paper include:

- **Aromatic levels in market fuels decreased by approximately 7% by volume for each 10% by volume increase in ethanol content** (Table 1). Our findings of lower aromatic content with increasing ethanol content is consistent with market fuel studies by EPA and others (Eastern Research Group, 2017, Eastern Research Group, 2020, US EPA, 2017). As discussed in EPA's Fuel Trends Report, for example, ethanol volume in market fuels increased by approximately 9.4% between 2006 and 2016, while aromatics over the same time period were found to drop by 5.7% (US EPA, 2017).

We note that our estimated market fuel properties differ from those used in the recent US EPA Anti-Backsliding Study (ABS), which examined the impacts of changes in vehicle and engine emissions from ethanol-blended fuels on air quality (US EPA, 2020). Contrary to our study, ABS was based on hypothetical fuels that were intended to satisfy experimental considerations rather than mimic real-world fuels. It did not consider published fuel trends; rather, the ABS used inaccurate fuel property adjustment factors in its modeling, reducing aromatics by only 2% (Table 5.3 of ABS 2020), substantially lower than the reductions found in our paper and in fuel survey data (Kazemiparkouhi et al., 2021, US EPA, 2017). As a result, the ABS's findings and their extension to public health impacts are not generalizable to real world conditions.

Table 1. Estimated market fuel properties

| Fuel ID | EtOH Vol (%) | T50 (°F) | T90 (°F) | Aromatics Vol (%) | AKI | RVP (psi) |
|--|--------------|----------|----------|-------------------|-----|-----------|
| E0 | 0 | 219 | 325 | 30 | 87 | 8.6 |
| E10 | 10 | 192 | 320 | 22 | 87 | 8.6 |
| E15 | 15 | 162 | 316 | 19 | 87 | 8.6 |
| E20 | 20 | 165 | 314 | 15 | 87 | 8.6 |
| E30 | 30 | 167 | 310 | 8 | 87 | 8.6 |
| Abbreviations: EtOH = ethanol volume; T50 = 50% volume distillation temperature; T90 = 90% volume distillation temperature; Aromatics=aromatic volume; AKI = Anti-knock Index; RVP = Reid Vapor Pressure. | | | | | | |


- **PM emissions decreased with increasing ethanol content under cold-start conditions.** Primary PM emissions decreased by 15-19% on average for each 10% increase in ethanol content under cold-start conditions (Figure 1). While statistically significant for both engine types, PM emission reductions were larger for GDI as compared to PFI engines, with 53% and 29% lower PM emissions, respectively, when these engines burned E30 as compared to E10. In contrast, ethanol content in market fuels had no association with PM emissions during hot-running conditions.

Importantly, our findings are consistent with recent studies that examined the effect of ethanol blending on light duty vehicle PM emissions. Karavalakis et al. (2014), (2015), Yang et al. (2019a), (2019b), Schuchmann and Crawford (2019), for

example, assessed the influence of different mid-level ethanol blends – with proper adjustment for aromatics – on the PM emissions from GDI engines and Jimenez and Buckingham (2014) from PFI engines. As in our study, which also adjusted for aromatics, each of these recent studies found higher ethanol blends to emit lower PM as compared to lower or zero ethanol fuels.

Together with these previous studies, our findings support the ability of ethanol-blended fuels to offer important PM emission reduction opportunities. **Cold start PM emissions have consistently been shown to account for a substantial portion of all direct tailpipe PM emissions from motor vehicles**, with data from the EPA study estimating this portion to equal 42% (Darlington et al., 2016, US EPA, 2013). The cold start contribution to total PM vehicle emissions, together with our findings of emission reductions during cold starts, suggest that a **10% increase in ethanol fuel content from E10 to E20 would reduce total tailpipe PM emissions from motor vehicles by 6-8%.**

Figure 1. Change (%) in cold-start emissions for comparisons of different ethanol-content market fuels^a



^a Emissions were predicted from regression models that included ethanol and aromatics volume fraction, T90, and RVP as independent variables

- **NO_x, CO and THC emissions were significantly lower for higher ethanol fuels for PFI engines under cold-start conditions**, but showed no association for GDI engines (Figure 1). CO and THC emissions also decreased under hot running conditions for PFI and for CO also for GDI engines (results not shown). [Note that NO_x emissions for both PFI and GDI engines were statistically similar for comparisons of all ethanol fuels, as were THC emissions for GDI engines.] These findings add to the scientific evidence demonstrating emission reduction benefits of ethanol fuels for PM and other key motor vehicle-related gaseous pollutants.

Implications for Public Health and Environmental Justice Communities

The estimated reductions in air pollutant emissions, particularly of PM and NO_x, indicate that increasing ethanol content offers opportunities to improve air quality and public health. As has been shown in numerous studies, lower PM emissions result in lower ambient PM concentrations and exposures (Kheirbek et al., 2016, Pan et al., 2019), which, in turn, are causally associated with lower risks of total mortality and cardiovascular effects (Laden et al., 2006, Pun et al., 2017, US EPA, 2019, Wang et al., 2020).

The above benefits to air quality and public health associated with higher ethanol fuels may be particularly great for environmental justice (EJ) communities. EJ communities are predominantly located in urban neighborhoods with high traffic density and congestion and are thus exposed to disproportionately higher concentrations of PM emitted from motor vehicle tailpipes (Bell and Ebisu, 2012, Clark et al., 2014, Tian et al., 2013). Further, vehicle trips within urban EJ communities tend to be short in duration and distance, with approximately 50% of all trips in dense urban communities under three miles long (de Nazelle et al., 2010, Reiter and Kockelman, 2016, US DOT, 2010). As a result, a large proportion of urban vehicle trips occur under cold start conditions (de Nazelle et al., 2010), when PM emissions are highest. Given the evidence that ethanol-blended fuels substantially reduce PM, NO_x, CO, and THC emissions during cold-start conditions, it follows that ethanol-blended fuels may represent an effective method to reduce PM health risks for EJ communities.

Summary

Findings from Kazemiparkouhi et al. (2021) provide important, new evidence of ethanol-related reductions in vehicular emissions of PM, NO_x, CO, and THC based on real-world fuels and cold-start conditions. Given the substantial magnitude of these reductions and their potential to improve air quality and through this public health, our findings warrant careful consideration. Policies that encourage higher concentrations of ethanol in gasoline would provide this additional benefit. These policies are especially needed to protect the health of EJ communities, who experience higher exposures to motor vehicle pollution, likely including emissions from cold starts in particular, and are at greatest risk from their effects.

References

- BELL, M. L. & EBISU, K. 2012. Environmental inequality in exposures to airborne particulate matter components in the United States. *Environmental health perspectives*, 120, 1699-1704.
- CLARK, L. P., MILLET, D. B. & MARSHALL, J. D. 2014. National patterns in environmental injustice and inequality: outdoor NO₂ air pollution in the United States. *PLoS One*, 9, e94431.
- DARLINGTON, T. L., KAHLBAUM, D., VAN HULZEN, S. & FUREY, R. L. 2016. Analysis of EPA Act Emission Data Using T70 as an Additional Predictor of PM Emissions from Tier 2 Gasoline Vehicles. *SAE Technical Paper*.
- DE NAZELLE, A., MORTON, B. J., JERRETT, M. & CRAWFORD-BROWN, D. 2010. Short trips: An opportunity for reducing mobile-source emissions? *Transportation Research Part D: Transport and Environment*, 15, 451-457.
- EASTERN RESEARCH GROUP 2017. Summer Fuel Field Study (prepared for Texas Commission on Environmental Quality by Eastern Research Group, Inc.).
- EASTERN RESEARCH GROUP 2020. Summer Field Study (prepared for Texas Commission on Environmental Quality by Eastern Research Group, Inc.).
- JIMENEZ, E. & BUCKINGHAM, J. P. 2014. Exhaust Emissions of Average Fuel Composition. Alpharetta, GA.
- KARAVALAKIS, G., SHORT, D., VU, D., RUSSELL, R. L., ASA-AWUKU, A., JUNG, H., JOHNSON, K. C. & DURBIN, T. D. 2015. The impact of ethanol and iso-butanol blends on gaseous and particulate emissions from two passenger cars equipped with spray-guided and wall-guided direct injection SI (spark ignition) engines. *Energy*, 82, 168-179.
- KARAVALAKIS, G., SHORT, D., VU, D., VILLELA, M., ASA-AWUKU, A. & DURBIN, T. D. 2014. Evaluating the regulated emissions, air toxics, ultrafine particles, and black carbon from SI-PFI and SI-DI vehicles operating on different ethanol and iso-butanol blends. *Fuel*, 128, 410-421.
- KAZEMIPARKOUHI, F., ALARCON FALCONI, T. M., MACINTOSH, D. L. & CLARK, N. 2021. Comprehensive US database and model for ethanol blend effects on regulated tailpipe emissions. *Sci Total Environ*, 151426.
- KHEIRBEK, I., HANEY, J., DOUGLAS, S., ITO, K. & MATTE, T. 2016. The contribution of motor vehicle emissions to ambient fine particulate matter public health impacts in New York City: a health burden assessment. *Environmental Health*, 15, 89.
- LADEN, F., SCHWARTZ, J., SPEIZER, F. E. & DOCKERY, D. W. 2006. Reduction in fine particulate air pollution and mortality: Extended follow-up of the Harvard Six Cities study. *American journal of respiratory and critical care medicine*, 173, 667-672.
- PAN, S., ROY, A., CHOI, Y., ESLAMI, E., THOMAS, S., JIANG, X. & GAO, H. O. 2019. Potential impacts of electric vehicles on air quality and health endpoints in the Greater Houston Area in 2040. *Atmospheric Environment*, 207, 38-51.
- PUN, V. C., KAZEMIPARKOUHI, F., MANJOURIDES, J. & SUH, H. H. 2017. Long-Term PM_{2.5} Exposure and Respiratory, Cancer, and Cardiovascular Mortality in Older US Adults. *American Journal of Epidemiology*, 186, 961-969.
- REITER, M. S. & KOCKELMAN, K. M. 2016. The problem of cold starts: A closer look at mobile source emissions levels. *Transportation Research Part D: Transport and Environment*, 43, 123-132.

- SCHUCHMANN, B. & CRAWFORD, R. 2019. Alternative Oxygenate Effects on Emissions. Alpharetta, GA (United States).
- TIAN, N., XUE, J. & BARZYK, T. M. 2013. Evaluating socioeconomic and racial differences in traffic-related metrics in the United States using a GIS approach. *J Expo Sci Environ Epidemiol*, 23, 215-22.
- US DOT 2010. National Transportation Statistics. Research and Innovative Technology Administration: Bureau of Transportation Statistics.
- US EPA 2013. Assessing the Effect of Five Gasoline Properties on Exhaust Emissions from Light-Duty Vehicles Certified to Tier 2 Standards: Analysis of Data from EPAAct Phase 3 (EPAAct/V2/E-89): Final Report. EPA-420-R-13-002 ed.: Assessment and Standards Division Office of Transportation and Air Quality U.S. Environmental Protection Agency.
- US EPA 2017. Fuel Trends Report: Gasoline 2006-2016.
- US EPA 2019. Integrated Science Assessment for Particulate Matter. Center for Public Health and Environmental Assessment.
- US EPA 2020. Clean Air Act Section 211(v)(1) Anti-backsliding Study. Assessment and Standards Division Office of Transportation and Air Quality U.S. Environmental Protection Agency.
- WANG, B., EUM, K. D., KAZEMIPARKOUHI, F., LI, C., MANJOURIDES, J., PAVLU, V. & SUH, H. 2020. The impact of long-term PM_{2.5} exposure on specific causes of death: exposure-response curves and effect modification among 53 million U.S. Medicare beneficiaries. *Environ Health*, 19, 20.
- YANG, J., ROTH, P., DURBIN, T. D., JOHNSON, K. C., ASA-AWUKU, A., COCKER, D. R. & KARAVALAKIS, G. 2019a. Investigation of the Effect of Mid- And High-Level Ethanol Blends on the Particulate and the Mobile Source Air Toxic Emissions from a Gasoline Direct Injection Flex Fuel Vehicle. *Energy & Fuels*, 33, 429-440.
- YANG, J., ROTH, P., ZHU, H., DURBIN, T. D. & KARAVALAKIS, G. 2019b. Impacts of gasoline aromatic and ethanol levels on the emissions from GDI vehicles: Part 2. Influence on particulate matter, black carbon, and nanoparticle emissions. *Fuel*, 252, 812-820.